

Ecological correlates of *ex situ* seed longevity: a comparative study on 195 species

Robin J. Probert*, Matthew I. Daws† and Fiona R. Hay

Seed Conservation Department, Royal Botanic Gardens Kew, Wakehurst Place, Ardingly, West Sussex RH17 6TN, UK

Received: 1 January 2009 Returned for revision: 17 February 2009 Accepted: 9 March 2009 Published electronically: 9 April 2009

- **Background and Aims** Extended seed longevity in the dry state is the basis for the *ex situ* conservation of ‘orthodox’ seeds. However, even under identical storage conditions there is wide variation in seed life-span between species. Here, the effects of seed traits and environmental conditions at the site of collection on seed longevity is explored for 195 wild species from 71 families from environments ranging from cold deserts to tropical forests.
- **Methods** Seeds were rapidly aged at elevated temperature and relative humidity (either 45°C and 60% RH or 60°C and 60% RH) and regularly sampled for germination. The time taken in storage for viability to fall to 50% (p_{50}) was determined using Probit analysis and used as a measure of relative seed longevity between species.
- **Key Results** Across species, p_{50} at 45°C and 60% RH varied from 0.1 d to 771 d. Endospermic seeds were, in general, shorter lived than non-endospermic seeds and seeds from hot, dry environments were longer lived than those from cool, wet conditions. These relationships remained significant when controlling for the effects of phylogenetic relatedness using phylogenetically independent contrasts. Seed mass and oil content were not correlated with p_{50} .
- **Conclusions** The data suggest that the endospermic seeds of early angiosperms which evolved in forest understorey habitats are short-lived. Extended longevity presumably evolved as a response to climatic change or the invasion of drier areas. The apparent short-lived nature of endospermic seeds from cool wet environments may have implications for re-collection and re-testing strategies in *ex situ* conservation.

Key words: Gene banks, seed ageing, seed longevity, taxonomic trends, climate, seed structure.

INTRODUCTION

The fact that seeds of most species can be dried and stored from year-to-year has been exploited since the beginning of agriculture. Indeed, the ability of many orthodox seeds (*sensu* Roberts, 1973) to remain viable for tens or hundreds of years in dry storage (Walters *et al.*, 2005; Daws *et al.*, 2007), means that they also provide a convenient vehicle for the long-term *ex situ* conservation of plant germplasm.

The longevity of seeds held in dry storage is mainly determined by seed moisture content and storage temperature, with life-span increasing predictably with decreasing temperature and moisture content (Harrington, 1963, 1972; Ellis and Roberts, 1980). However, there are also wide inherent differences in seed longevity between species (Harrington, 1972; Priestley *et al.*, 1985). For example, using the improved seed viability equations (derived from rapid ageing experiments at elevated temperatures and moisture contents; Ellis and Roberts, 1980), the predicted time for viability to decline from 97.7% to 84.1% for seeds stored under gene-bank conditions (–20°C after equilibration at 15% RH, 15°C) ranges from approx. 30 years for *Ulmus carpinifolia* to approx. 6000 years for *Sorghum bicolor* (Liu *et al.*, 2008). Similarly, using the Avrami equation to model data from re-testing of 41 847 seed accessions from 276 species stored at the USDA National Center for Genetic Resources Preservation (NCGRP), Walters *et al.* (2005) predicted that there would be a difference of 626 years in the time for viability to fall

to 50% between the shortest (*Bromus stichensis*; 7 years) and longest lived accessions (*Trifolium campestre*; 633 years).

However, there is already evidence that some species produce seeds with much shorter longevity in dry storage. For example, seeds (with high initial viability) of *Anemone nemorosa* are predicted to survive <1 year under seed bank storage conditions (Ali *et al.*, 2007). As there is an increasing aspiration to conserve seeds from wild plant species (Target VIII of the Global Strategy for Plant Conservation; SCBD, 2006), it is likely that more species will be found whose seeds have a similarly short life-span.

Understanding species differences in seed longevity is therefore crucial to the effective management of seed conservation collections because it underpins the selection of viability re-test intervals and hence regeneration or re-collection strategies. It is particularly critical for collections of wild plant species where, due to the genetic heterogeneity of wild plant populations, any significant decline in viability will result in the loss of genotypes from the accession (Walters, 2003). However, species-specific constants for the improved viability equations are only available for approx. 56 species (Liu *et al.*, 2008) the majority of which are temperate species of agricultural importance. Generating constants is time consuming and involves the expenditure of tens-of-thousands of seeds per species. Even if the universal values for the temperature coefficients (Dickie *et al.*, 1990) are accepted, carrying out the rapid ageing experiments needed to determine the other two parameters would still require thousands of seeds. Thus, species constants are unlikely ever to be experimentally determined for the majority of plant species. Consequently, predictions of seed longevity in gene-bank storage for threatened

* For correspondence. E-mail r.probert@kew.org

† Present address: EWL Sciences Pty Ltd, GPO Box 518, Darwin, NT 0800, Australia.

wild plant species are problematic. An alternative approach to actually determining longevity for individual species is to identify reliable and robust correlates of longevity and use these to develop general predictive models (Daws *et al.*, 2006).

A number of studies have reported potential correlates of seed longevity in dry storage including seed mass, oil content, carbohydrate composition, taxonomy and climate (e.g. Horbowicz and Obendorf, 1994; Priestley, 1986; Pritchard and Dickie, 2003; Walters *et al.*, 2005). However, a purported link between high oil content and short storage life-span has not been supported by recent analyses (Pritchard and Dickie, 2003; Walters *et al.*, 2005; Bird, 2006).

Seed size may be used as a predictor of persistence in the natural environment, with small seeds generally showing greater persistence in the soil seed bank compared with larger seeds (Thompson *et al.*, 1993; Funes *et al.*, 1999; Cerabolini *et al.*, 2003; Peco *et al.*, 2003). It might be expected that seeds showing greater persistence in the soil would be long-lived in dry storage. This is supported by reports of a positive correlation between seed longevity measured in the laboratory under controlled (Long *et al.*, 2008) or uncontrolled (Bekker *et al.*, 2003) conditions and seed persistence in the soil. However, no such correlation was found when seed longevity under seed bank conditions was compared with seed persistence in soil (Walters *et al.*, 2005), possibly because the stresses and protective mechanisms involved in conferring persistence in the soil are different to the stresses and protective mechanisms that make seeds resistant to ageing in long-term dry storage at low temperatures.

The most significant advance in understanding the underlying factors influencing seed longevity in dry storage comes from the analysis of the NCGRP data. Walters *et al.* (2005) suggested both a taxonomic and climatic component to interspecific differences in longevity. Thus, seeds from some families were inherently short lived (e.g. Apiaceae, Brassicaceae) and others long-lived (e.g. Malvaceae and Chenopodiaceae) and species originating from cool, temperate climates tended to produce seeds with short life-spans and those from warm, arid environments seeds with long life-spans. However, these authors did not assess the relative importance of these factors.

In this paper is presented a predictive model of relative seed longevity, determined by analysing longevity data for 195 species sampled from accessions stored in the Royal Botanic Gardens (RBG) Kew, Millennium Seed Bank and representing diverse taxa and global-wide sampling. Due to the logistical and time-scale problems with comparing storage life-span under gene-bank conditions, seed longevity was determined using standardized rapid ageing conditions (Davies and Probert, 2004; Hay *et al.*, 2006). These data are used to assess putative factors of significance in predicting seed longevity, including physical attributes of the seed such as mass and oil content and the climate of the area where seed collections originated. Relative embryo size (RES) across species has also been correlated with the developmental stage of the embryo (Forbis *et al.*, 2002) and intra-specific studies have indicated that longevity increases during seed development (e.g. Hay and Probert, 1995; Ali *et al.*, 2007). Therefore, we also consider whether differences in the relative size of the embryo are related to differences in storage life-span. In

addition, since performance at sub-zero temperatures has to be extrapolated from rapid ageing conditions, we test whether our ageing data correlates with long-term (20-year) storage data for species held under gene-bank conditions.

Specifically, the following hypotheses are tested with the aim of developing a predictive model for seed longevity: (1) among seed traits, seed mass, oil content and RES will not be related to seed longevity; (2) climatic parameters (mean annual temperature and total annual rainfall) will not be related to seed longevity. In comparative analyses, species are not statistically independent, as demonstrated by Walters *et al.* (2005) for seed longevity. Therefore, not only will cross-species analyses be conducted but also phylogenetically independent contrasts will be conducted to test for an evolutionary association between the variables of interest and seed storage life-span.

MATERIALS AND METHODS

Species selection

High viability collections ($\geq 85\%$ germination) of 195 species with known germination requirements were identified from conservation collections held at the Millennium Seed Bank under international gene bank standard conditions of 15% RH and -20°C (FAO/IPGRI, 1994) for between 2 and 35 years (Table 1). Species were selected to give broad taxonomic and geographic coverage and were only included if seed longevity characteristics had not been studied previously. They represent a range of orders (30) and families (71), from 65°N (Finland) to 50°S (Chile) and 173°E (New Zealand) to 125°W (Canada).

Ageing protocol

Ageing experiments on samples from the targeted collections were carried out between 2001 and 2006. At the outset, all seed samples were aged according to a standardized protocol (Davies and Probert, 2004). First, in order to raise the moisture content of the seeds prior to ageing and to minimize the subsequent adjustment of moisture content when samples were transferred to the ageing conditions, ten samples of 50 seeds each were rehydrated at 47% RH at 20°C in open glass vials or Petri dishes; the vials/dishes were placed over a non-saturated solution of LiCl (anhydrous, Laboratory Reagent Grade; Fisher Scientific UK Ltd, Leics., UK) in distilled water held in a sealed $300 \times 300 \times 130$ mm electrical enclosure box (Ensto UK Ltd, Southampton). At the end of the re-hydration period (14 d), seed equilibrium relative humidity (eRH) was checked using a sample of the equilibrating seeds or, if the volume of seeds was too small, a reference sample of *Ranunculus sceleratus* seeds that accompanied the test species during rehydration. The eRH was measured using a water activity measuring instrument which comprised a hygrometer sensor housed in an AW-D10 water activity probe, used in conjunction with a HygroPalm 3 display unit (Rotronic Instruments UK Ltd, Crawley, UK). Once the test species was judged to have reached equilibrium, samples were transferred to a second electrical enclosure box, over a non-saturated solution of LiCl at 60% RH placed in a LEEC

KIF compact incubator (LEEC Ltd, Nottingham, UK) at $45 \pm 2^\circ\text{C}$. The RH generated by the LiCl in the box was checked at 4- to 6-week intervals by pipetting a sample of approx. 10 mL solution and placing it into the sample chamber of the water activity-measuring instrument described above. Samples were allowed to equilibrate for up to 2 h before returning the sample to the enclosure box. The bulk solution was adjusted if necessary, usually by adding distilled water, stirring and allowing the solution to equilibrate before rechecking the RH (Hay *et al.*, 2008).

One sample of 50 seeds for each species was removed after each of 1, 2, 5, 9, 20, 30, 50, 75, 100 and 125 d for germination testing. Seeds were sown as two lots of 25 seeds each on 1% distilled water agar held in 55- or 90-mm-diameter Petri dishes (Bibby Sterilin Ltd, Staffordshire, UK) and placed in a LMS 250A or 600A cooled incubator (LMS Ltd, Sevenoaks, UK) at a temperature regime (constant or alternating) previously found to be optimal for germination of that accession in routine seed bank testing (see Appendix). Some of the accessions required additional treatments including the addition of 250 mg L^{-1} GA₃ (Sigma-Aldrich Company Ltd, Dorset, UK), mechanical scarification of the seed coat (carried out prior to equilibration at 49% RH) or cold stratification (typically at 5°C for 56 d) (Appendix). Plates were regularly checked for germination and seeds scored as germinated once the radicle had reached ≥ 2 mm in length.

In the ageing experiments at 60% RH and 45°C , four species of Myrtaceae: *Calothamnus crassus*, *C. graniticus*, *C. rupestris* and *Melaleuca diosmifolia* were found to be extremely long-lived, requiring sampling up to 500 d to generate reliable survival curves. In order to speed up the ageing process for a further 20 species, which were also expected to be very long lived, ageing was carried out at 60% RH and 60°C . These species originated mainly from Australia and/or displayed serotiny. Sampling was carried out at 2, 5, 10, 15, 20, 25, 30, 40, 55 and 75 d and a germination test carried out as before.

Determination of seed characteristics

The mean seed air dry weight (seeds equilibrated to 15% RH, 15°C) was calculated by weighing five lots of 50 seeds for each accession used in the study. Where available, whole seed oil contents for the species were obtained from either the Seed Information Database of the Royal Botanic Gardens Kew (Liu *et al.*, 2008) or the Seed Oils Fatty Acid Database (<http://sofa.bfel.de/>).

RES was determined following the method of Forbis *et al.*, (2002). Five seeds from each targeted accession were sectioned longitudinally. Subsequently, digital images were taken using a Stemi SV 11 Microscope (Carl Zeiss, Germany) and using Axiovision 3.1 (Carl Zeiss Vision), the area of the embryo was measured and expressed as a proportion of the total space inside the testa. However, since RES was skewed towards values close to 1.0 and could not be satisfactorily transformed to approximate the normal distribution, a discrete classification was used in all analyses where seeds were classified as endospermic when the RES is < 1 and non-endospermic when the RES = 1.

Estimation of climate for seed-lot collection sites

Annual mean temperature and total annual precipitation values, based on data collated between 1950 and 2000, were obtained for the collection location of each seed lot by querying WORLDCLIM data (download version 1.4 <http://www.worldclim.org/>) at a maximum resolution of 30 arc-seconds (approx. 1 km) using the 'Extract Values to Points' tool in ESRI ArcMap (version 9.1).

Seed bank data analysis

For seed lots which have been held for > 20 years at the Millennium Seed Bank, data for initial germination and germination after 20 years were extracted from the Millennium Seed Bank's Seed Bank Database. The highest percentage germination result in initial tests and after 20 years' storage were compared to test whether there had been a significant drop in viability during storage. Collections in which there had been a significant drop in germinability were identified by testing the null hypothesis of no difference between the two germination values. This was achieved by calculating the two-tailed probability corresponding to Z , the value of the normal deviate corresponding to the difference between the initial and final germination percentage values following the procedure outlined in Ellis *et al.* (1985).

As a result of limited overlap between both species and genera in this 20-year analysis and species in the artificial ageing tests, the proportion of species, showing a significant decline in viability over 20 years was calculated at the family level. This value was then correlated with the mean p_{50} for corresponding families (from rapid ageing) to test for a relationship between survival life-span under gene-bank and rapid ageing conditions.

Statistical analysis

Probit analysis of the seed ageing data was carried out using GenStat for Windows, Version 8 (VSN International Ltd, Oxford, UK) to estimate p_{50} (the time for viability to fall to 50%) through fitting of the basic seed viability equation:

$$v = K_i - (p/\sigma) \quad (1)$$

where v is the viability in normal equivalent deviates (NED) at time p (days); K_i is the initial viability (NED) and σ is the standard deviation of the normal distribution of seed deaths in time.

For seeds aged at the higher temperature (60°C), an estimate of p_{50} at 45°C was calculated by applying a correction factor based on data for five species tested at both temperatures (three species tested in our laboratory and two species tested by collaborators, A. Martyn and D. Merritt). Across these five species there was an average 10.9 (\pm s.d. 5.7) fold difference in p_{50} between the two temperatures. Whilst the corrected values must be treated with caution, estimates of longevity (time for viability to fall by 1 NED) under the two different ageing environments calculated using the Seed Information Database (Liu *et al.*, 2008), for seeds of other species for which viability constants for the Ellis and Roberts (1980)

TABLE 1. Comparison between the proportion of seed collections per family displaying a significant ($P < 0.05$) drop in viability during 20-year storage at 15% RH and -20°C in the RBG Kew, Millennium Seed Bank, and the mean p_{50} recorded for that family from artificial ageing at 45°C and 60% RH (also shown are the ranks used for performance at -20°C and in artificial ageing that were used in the Spearman's Rank Correlation)

Family (-aceae)	No. of collections tested after 20 years	Proportion of collections displaying no significant drop in germination after 20 years	Rank 1	Mean p_{50} (d)	n	Rank 2
Aizo-	2	0.50	16	76.9	1	42
Amaranth-	64	0.84	41	38.7	4	22
Api-	29	0.55	20	54.4	2	35
Aster-	207	0.75	36	52.6	4	31
Boragin-	8	0.50	16	47.0	1	27
Brassic-	73	0.70	29	43.6	2	26
Campanul-	32	0.41	12	11.2	4	11
Caprifoli-	1	0.00	1	8.3	1	7
Caryophyll-	513	0.81	40	39.8	16	24
Cercidiphyll-	1	0.00	1	0.1	1	1
Cist-	30	0.93	46	78.2	1	43
Convolvul-	10	0.90	45	69.3	2	41
Crassul-	22	0.73	33	57.0	2	36
Cucurbit-	7	0.86	42	54.0	1	34
Cupress-	1	0.00	1	10.7	1	10
Cyper-	8	0.75	35	47.0	2	28
Dipsac-	9	0.78	38	59.3	3	38
Eric-	13	0.31	9	9.4	5	9
Fab-	5	0.76	37	83.5	6	45
Gentian-	51	0.71	32	19.2	6	15
Gerani-	5	0.80	39	82.8	2	44
Gesneri-	1	0.00	1	0.9	1	2
Haemodor-	2	0.50	16	41.7	1	25
Hydrophyll-	5	0.40	11	49.5	2	29
Irid-	9	0.67	26	53.7	1	32
Junc-	8	0.63	22	59.2	1	37
Juncagin-	1	0.00	1	5.6	1	5
Lami-	75	0.65	25	69.0	7	40
Loas-	5	0.60	21	9.0	2	8
Lythr-	3	0.67	26	14.0	1	12
Melanthi-	1	0.00	1	4.5	4	3
Onagr-	171	0.49	15	53.9	12	33
Papaver-	49	0.45	13	36.7	3	21
Po-	722	0.86	44	26.7	4	19
Polemoni-	4	0.50	16	22.0	3	17
Polygon-	20	0.70	30	38.7	1	23
Portulac-	11	0.73	33	51.1	1	30
Primul-	30	0.47	14	31.0	8	20
Ranuncul-	81	0.69	28	14.9	6	13
Ros-	56	0.86	42	21.8	2	16
Rubi-	10	0.70	30	16.0	3	14
Saxifrag-	3	0.33	10	4.6	2	4
Scrophulari-	101	0.63	24	25.4	4	18
Solan-	75	0.63	23	62.9	8	39
Urtic-	2	0.00	1	84.4	1	46
Viol-	2	0.00	1	6.3	2	6

viability equations have been determined and for which seed oil contents are known (to estimate moisture contents under the two different ageing environments), suggest that this correction factor is appropriate.

Oil content was only available for 140 of 195 species. Consequently two analyses were conducted for both the multiple regression and phylogenetically independent contrasts (PIC) approaches. The first analysis included only the species for which oil content was available, the second all 195 species but excluding the oil content data. However, since oil content was the first term to be dropped from the regression model for the 140 species (i.e. it contributed

least to explaining variation in p_{50}) in both the cross-species and PIC approaches, only the multiple regression results for the analyses involving all 195 species are presented here. In the multiple regression analyses, the relationship between seed longevity (p_{50}) and oil content (% dry weight basis), seed mass (mg), presence (1)/absence of endosperm (0), mean annual temperature ($^{\circ}\text{C}$) and total annual rainfall (mm) was analysed using Minitab version 13 (Minitab Inc., State College, PA, USA). To ensure normality, p_{50} and seed mass were \log_{10} transformed and oil content was arc-sine transformed in this and all other analyses. Total annual rainfall was transformed using a Box-Cox

transformation with the optimum value of λ (0.30) determined using Minitab 13.

Regression models were constructed using backwards elimination. Initially all terms were included in the model and at each step in the procedure the variable with the smallest (non-significant) partial correlation was dropped from the model until only significant ($P < 0.05$) terms remained in the model (Sokal and Rohlf, 1995).

PICs (Felsenstein, 1985; Pagel, 1992) were used to analyse the relationship between seed longevity (as assessed by p_{50}) and seed mass, oil content, the occurrence of endosperm, mean annual temperature and total annual rainfall. This approach is based on the logic of comparing pairs of species within a phylogeny that share an immediate common ancestor. The null hypothesis is that there is no correlation between changes in traits at the nodes. The package CAIC (Purvis and Rambaut, 1995) was used to generate contrasts. Within the phylogeny it was assumed that all branch lengths were the same: analyses of simulated data sets suggest that equal branch lengths may perform better than estimated branch lengths (Purvis *et al.*, 1994). For analyses testing whether the transition from endospermic to non-endospermic seeds (or vice versa) is associated with changes in p_{50} , the Brunch procedure, designed for discrete predictor variables, was used (Purvis and Rambaut, 1995). Contrasts from Brunch were analysed using a *t*-test on the mean of the contrasts: a mean significantly different to zero indicates correlated evolution between the traits of interest. For the continuous variables (seed mass, oil content, etc.) the Crunch procedure was used (Purvis and Rambaut, 1995). For Crunch, variables were either run individually, with a linear regression forced through the origin, subsequently fitted to the contrasts or simultaneously with multiple regression used to analyse the contrasts. For all phylogenetic analyses, the latest phylogeny available to (sub-) family level from the Angiosperm Phylogeny Group was used (APG II, 2003). However, due to the wide range of families and the lack of complete

phylogenies to genus level for most families, a series of polytomies were created.

RESULTS

Variation in p_{50} between species

Across the 195 species, viability declined with increasing duration of the ageing treatment (for examples, see Fig. 1). However, there was wide variation across species in the time taken for viability to fall to 50%: p_{50} ranged between 0.1 d for *Cercidiphyllum japonicum* to 771 d for *Calothamnus rupestris*, respectively (both aged at 45°C and 60% RH; Appendix). Within orders, there was a wide span of longevities with p_{50} spanning more than an order of magnitude in some orders (e.g. Fagales and Lamiales). In addition, the Myrtales contained the longest lived species and the Liliales the shortest lived (Fig. 2A). Similarly there was wide variation in p_{50} between families, including in some cases within the same order (e.g. Myrtaceae and Onagraceae in the Myrtales with mean estimates for p_{50} of 366 d and 54 d, respectively; Fig. 2B).

For the 46 families that overlapped between rapid ageing and 20 years' seed bank storage, there was a highly significant correlation between the proportion of species in a family exhibiting a significant decline in viability after 20 years storage under gene-bank conditions and mean p_{50} at the family level (Spearman's Rank Correlation, $r_s = 0.527$, d.f. = 44, $P < 0.001$; Table 1).

Correlates of p_{50}

Endospermic seeds were significantly shorter lived than non-endospermic seeds (mean p_{50} values of 20.3 d vs. 65.7 d, one-way ANOVA, $P < 0.001$; Fig. 3). However, neither of the additional seed characteristics (seed mass and oil content) was significantly related to p_{50} (linear regression,




FIG. 1. Survival curves fitted by probit analysis for four of the species in this study: seeds of (A) *Tofieldia pusilla*, *Blackstonia perfoliata* and *Calothamnus crassus* aged at 60% RH and 45°C, and (B) *Calothamnus rupestris* aged at 60% RH, 60°C and 60% RH, 45°C (inset).




FIG. 2. Box plots of p_{50} in (A) orders and (B) families. Only orders and families with three or more species tested are shown. Boxes span the 25th to 75th percentiles; whiskers span the 5th to 95th percentiles.




FIG. 3. Box plots comparing p_{50} for endospermic and non-endospermic seeds. Boxes span the 25th to 75th percentiles; whiskers span the 5th to 95th percentiles.

$R^2 \leq 0.02$, $P > 0.05$; Fig. 4A, B). There was a significant positive relationship between mean annual temperature and p_{50} (linear regression, $R^2 = 0.102$, $P < 0.001$), species from warmer environments tending to have a greater p_{50} (Fig. 4C). There was also a significant negative relationship between total annual precipitation (Box-Cox transformed data, $\lambda = 0.30$) and p_{50} (linear regression, $R^2 = 0.051$, $P = 0.002$), species from drier environments tending to have a greater p_{50} (Fig. 4D).

Multiple regression for all 195 species, including the presence/absence of endosperm, \log_{10} (seed mass), mean annual temperature and total annual precipitation (Box-Cox transformed, $\lambda = 0.30$) as the independent variables and p_{50} as the dependent variable, explained a significant proportion of the variation in p_{50} ($R^2 = 0.279$, $F = 18.39$, $P < 0.001$). Only the presence/absence of endosperm, mean annual temperature and total rainfall remained when using a backwards model selection procedure (Table 2). This model explained a significant proportion of the variation in time for viability to

fall to 50% ($R^2 = 0.278$, $F = 24.5$, $P < 0.001$) such that:

$$\log_{10}(p_{50}) = 2.06 - 0.459(E) + 0.021(AT) - 0.081(AR^{0.3})$$

where p_{50} is the time for viability to fall to 50% (days), E is the presence (1) or absence (0) of endosperm, AT is mean annual temperature ($^{\circ}\text{C}$) and AR is annual rainfall (mm).

Accounting for phylogenetic relatedness indicated that the presence of endosperm was significantly associated with lower values of p_{50} (1-sample t -test, $t = -2.36$, $P < 0.05$). There was no relationship between contrasts in oil content and p_{50} ($P > 0.05$, Fig. 5A) or contrasts in seed mass and p_{50} ($P > 0.05$, Fig. 5B). However, there were significant relationships between contrasts in temperature and p_{50} (positive) and rainfall and p_{50} (negative) ($P < 0.05$; Fig. 5C, D). Simultaneously generating contrasts for all three continuous variables (excluding oil content) and p_{50} in Crunch and then conducting multiple regression on the contrasts showed a significant relationship ($F = 2.70$, $P < 0.05$). Partial correlation showed a significant negative correlation between p_{50} and annual rainfall contrasts ($\beta = -0.286$, $t = -2.29$, $P < 0.05$) and no correlation between p_{50} and either seed mass ($\beta = 0.029$, $t = 0.4$, $P = 0.687$) or mean annual temperature ($\beta = 0.01$, $t = 1.03$, $P = 0.304$).

DISCUSSION

The present data indicate variation in seed longevity spanning four orders of magnitude under identical storage conditions. This variation was related to seed structure and the climate of the area from where the species were collected. In addition, it was found that survival of rapid ageing, at the family level, was correlated with survival under gene-bank conditions. These findings may enable predictions of likely performance under gene-bank conditions based on information regarding climate and seed structure.




FIG. 4. Relationship between p_{50} and (A) oil content, (B) seed mass, (C) annual temperature, and (D) annual rainfall.

TABLE 2. Model selection for the effects of endosperm, seed mass, mean annual temperature and total annual rainfall on p_{50} in the multiple regression analysis

Term	Coefficient	t-value
Full model		
Constant	2.060	10.58***
Endosperm	-0.459	-5.86***
Mass	0.025	0.56 n.s.
Temperature	0.021	3.83***
Rainfall	-0.070	-3.36***
Reduced model		
Constant	2.059	10.59***
Endosperm	-0.459	-5.86***
Temperature	0.021	4.05***
Rainfall	-0.081	-3.51***

Terms in bold are those that were dropped from the model at each stage in the model selection process.

***, $P < 0.001$; n.s., not significant $P > 0.05$

Seed mass, lipid content and p_{50}

Among seed characteristics, it was found that, in agreement with other studies (Priestley, 1986; Walters *et al.*, 2005; Bird, 2006), there was no relationship between seed longevity and oil content. Nonetheless, it is possible that lipid composition is important in determining longevity since the degree of fatty acid saturation will impact on the potential for lipid peroxidation which has been considered to be a primary reaction in ageing contributing to free-radical production and

subsequent attack on other macro-molecules (Benson, 1990). This remains to be demonstrated.

Similarly, seed mass was not correlated with longevity as concluded elsewhere for a range of crop species (Pritchard and Dickie, 2003; Walters *et al.*, 2005). Previous reports have indicated that smaller seeds tend to persist longer in the natural environment (Thompson *et al.*, 1993; Funes *et al.*, 1999); however, this may be more to do with smaller seeds requiring particular environmental triggers for germination than greater longevity *per se*.

Walters *et al.* (2005) reported a lack of correlation between longevity under seed bank conditions and seed persistence in a temperate mesic soil, whereas Long *et al.* (2008) found a positive correlation between seed longevity under rapid ageing conditions (45°C and 60% RH) and seed persistence in soil. These apparently contradictory findings might be explained by the different soils investigated in the two studies. The comparatively warm dry soils used by Long and co-workers were certainly much closer to the conditions of the rapid ageing tests compared with the very different conditions of seed bank storage and temperate mesic soil investigated by Walters and co-workers.

Endosperm, environment, taxonomy and p_{50}

Whilst some of the variation in p_{50} is not explained by the present model, the analysis clearly demonstrates that species with endosperm (i.e. comparatively small embryos), from cool wet environments tend to be short-lived in dry storage.




FIG. 5. Relationship between p_{50} contrasts and (A) oil content contrasts, (B) seed mass contrasts, (C) annual temperature contrasts, (D) annual rainfall contrasts. Contrasts were generated in CAIC and the regressions were forced through the origin.

Moreover, these relationships are significant when accounting for the potentially confounding effects of phylogenetic relationships (using PICs). Similarly, Walters *et al.* (2005) observed that seeds of species that originated in moist, temperate geographic regions such as Europe tended to have shorter life-spans compared with seeds of species originating in warm arid regions (Australia, South Asia). In further support of the proposition that species from hot dry environments are more likely to have comparatively long-lived seeds, the survival and germination of 200-year-old seeds of three South African species has been recently reported (Daws *et al.*, 2007). Conversely, Ali *et al.* (2007) reported that endospermic seeds of *Anemone nemorosa* from the UK have very limited seed longevity (the time taken for viability to drop to 5% at 45°C and 60% RH ranged from approx. 2 d to 14 d): this species typically occurs in cool damp woodland environments.

In an evolutionary context, Forbis *et al.* (2002) attributed RES to the degree of embryo development at the time of dispersal with small embryos being the ancestral state in the angiosperms. Field and Arens (2005) suggested that early angiosperms evolved in disturbed, moist understorey environments such as montane forests and along riverbanks. Thus, the present data suggest that early angiosperms had short-lived seeds (in dry storage), which is likely to be consistent with their limited exposure to drying. Thus, we hypothesize that prolonged longevity in the dry state evolved as an adaptation to either climatic drying, or the invasion of hot, dry

environments where seeds might be expected to persist in the dry state for irregular multi-year intervals between rainfall events sufficient for successful establishment.

Figure 2 suggests that seed life-span is also a characteristic of particular taxonomic groupings, as reported previously (Priestley *et al.*, 1985; Walters *et al.*, 2005). For example, species investigated in the Myrtales were typically long-lived while those in the Liliales were short lived. Such patterns were also evident at the family level. For example, the Campanulaceae, Ericaceae and Melanthiaceae were consistently short lived (p_{50} ranged from 0.1 d to 22 d) whereas Myrtaceae were consistently long lived (p_{50} ranged from 152 d to 771 d). Within some families, p_{50} varied considerably, yet genera within those families showed relatively little variation in longevity. For example, in the Primulaceae, p_{50} ranged from 4 d to 140 d but the five species of *Primula* tested were all short lived (p_{50} ranged from 6 d to 22 d; cf. Fig. 2B and Appendix). All five species of *Gentiana* were also short lived (p_{50} ranged from 8 d to 13 d) whereas *Blackstonia perfoliata* and *Centaurium erythraea*, also Gentianaceae, were significantly longer lived (p_{50} of 38 d and 37 d, respectively). Hay *et al.* (2006) also found little variation in p_{50} within the genus *Rhododendron*. Thus, for these consistently short-lived genera, the longevity of one species is likely to be a good predictor of con-generics. However, these taxonomic trends may result from related species sharing similar traits such as seed structure, and habitats, and hence having a similar longevity. Interestingly, the

presence of endosperm and the environmental variables were still significantly related to p_{50} when these phylogenetic effects were controlled using PIC analysis, supporting the hypothesis that correlated evolution between p_{50} and other seed traits and climate has resulted in these apparent taxonomic trends.

Implications for gene banks

A significant challenge for seed-bank managers is ensuring collections maintain high levels of viability to avoid loss of genotypes from the population while storing seeds from increasingly diverse taxa, many of which have not been studied in any way (at least in terms of seed biology). Thus, predictions of seed longevity based on seed characteristics and species' ecology would be of significant benefit to the seed banking community. Current guidelines (Rao *et al.*, 2006) recommend that base collections stored according to international standards (FAO/IPGRI, 1994) be tested for viability after 5 or 10 years of storage depending on initial viability and whether the seeds have poor longevity. However, there is increasing evidence that seeds of some exceptionally short-lived species such as *Anemone nemorosa* may lose viability within a year or two (Ali *et al.*, 2007). Conversely, seeds that are likely to be extremely long lived in storage, such as some of the Myrtaceae species tested here, might currently be tested more frequently than necessary, involving the expenditure of seeds which could represent valuable germplasm (in the sense that limited numbers of seeds are available and recollection is difficult). Here, for the first time, data have highlighted that seed structure and climate of origin are important indicators of seed longevity. Thus species from cool, moist environments, particularly those with small embryos relative to the size of the seed, may need to be tested more frequently than non-endospermic seeds from hot dry environments.

In contrast to Walters *et al.* (2005), who investigated correlates of seed longevity under gene-bank conditions, seeds in the current study were aged at elevated temperatures and RH. Potential advantages of this approach include the application of a standardized protocol to a wide range of germplasm enabling rapid generation of data: this protocol was applied to 195 species from 30 orders and 71 families. However, while the causes of seed death in gene-bank and rapid-ageing conditions may not be the same, families with species whose seeds aged rapidly in artificial ageing also showed evidence of more rapid seed viability loss during seed bank storage, with a significant reduction in viability over 20 years storage at -20°C . Moreover, collections from genera, such as *Primula*, predicted to be consistently short lived in comparative ageing tests also showed significant declines in viability after 20 years' storage in the seed bank.

The present data have shown that seed longevity is related to seed structure and climate of origin. Short-lived seeds are more likely to possess small embryos and originate from cool wet regions. Although we have not been able to explain some of the variation in p_{50} and we have not considered intra-specific variation in seed longevity resulting from genetic or environmental influences or seed maturity, we

can now predict which species are more likely to be short lived in storage. The rapid-ageing test described here offers seed-bank managers a means of assessing the potential longevity of seed collections of such species under seed-bank conditions, thereby enabling the selection of appropriate viability retest intervals and thus better management of conservation collections.

ACKNOWLEDGEMENTS

We are grateful to Tiziana Ulian who helped with initial data analysis; Hannah Davies, Wendy Bastable, Kirsty Brazil and Rosemary Newton who carried out the controlled ageing tests and embryo measurements; Amelia Martyn and David Merritt for sharing data on two Australian species and many other Millennium Seed Bank partners who have taken an active interest and directly or indirectly supported this work. Financial support was provided by the Millennium Commission, The Wellcome Trust and Orange plc. The Royal Botanic Gardens, Kew is supported by grant-in-aid from Defra.

LITERATURE CITED

- Ali N, Probert R, Hay F, Davies H, Stuppy W. 2007. Post-dispersal embryo growth and acquisition of desiccation tolerance in *Anemone nemorosa* L. seeds. *Seed Science Research* **17**: 155–163.
- APG II. 2003. An update of the Angiosperm Phylogeny Group classification for the orders and families of flowering plants: APG II. *Botanical Journal of the Linnean Society* **141**: 399–436.
- Bekker RM, Bakker JP, Ozinga WA, Thompson K. 2003. Seed traits: essential for understanding seed longevity. *Aspects of Applied Biology* **69**: 1–9.
- Benson EE. 1990. *Free radical damage in stored germplasm*. Rome: International Board for Plant Genetic Resources.
- Bird AE. 2006. *Thermal methods and comparative seed longevity in the genus Ranunculus L.* PhD Thesis, University of London, UK.
- Cerabolini B, Ceriani RM, Caccianiga M, De Andreis R, Raimondi B. 2003. Seed size, shape and persistence in soil: a test on Italian flora from Alps to Mediterranean coasts. *Seed Science Research* **13**: 75–85.
- Davies H, Probert R. 2004. *Protocol for comparative seed longevity testing sheet*, www.kew.org/msbp/scitech/publications/comparative_longevity.pdf. London: Royal Botanic Gardens, Kew.
- Daws MI, Garwood NC, Pritchard HW. 2006. Prediction of desiccation sensitivity in seeds of woody species: a probabilistic model based on two seed traits and 104 species. *Annals of Botany* **97**: 667–674.
- Daws MI, Davies J, Vaes E, van Gelder R, Pritchard HW. 2007. Two-hundred-year seed survival of *Leucospermum* and two other woody species from the Cape Floristic region, South Africa. *Seed Science Research* **17**: 73–79.
- Dickie JB, Ellis RH, Kraak HL, Ryder K, Tompsett PB. 1990. Temperature and seed storage longevity. *Annals of Botany* **65**: 197–204.
- Ellis RH, Roberts EH. 1980. Improved equations for the prediction of seed longevity. *Annals of Botany* **45**: 13–30.
- Ellis RH, Hong TD, Roberts EH. 1985. *Handbook of seed technology for genebanks*. Vol. I. *Principles and methodology*. Rome: International Board for Plant Genetic Resources.
- FAO/IPGRI. 1994. *Genebank standards*. Rome: Food and Agriculture Organisation of the United Nations/International Plant Genetic Resource Institute.
- Felsenstein J. 1985. Phylogenies and the comparative method. *American Naturalist* **125**: 1–15.
- Field TS, Arens NC. 2005. Form, function and environments of the early angiosperms: merging extant phylogeny and ecophysiology with fossils. *New Phytologist* **166**: 383–408.
- Forbis TA, Floyd SK, de Queiroz A. 2002. The evolution of embryo size in angiosperms and other seed plants: implications for the evolution of seed dormancy. *Evolution* **56**: 2112–2125.

- Funes G, Basconcelo S, Diaz S, Cabido M. 1999. Seed size and shape are good predictors of seed persistence in soil in temperate mountain grasslands of Argentina. *Seed Science Research* 9: 341–345.
- Harrington JF. 1963. The value of moisture-resistant containers in vegetable seed packaging. *Californian Agricultural Station Bulletin* 792: 1–23.
- Harrington JF. 1972. Seed storage and longevity. In: Kozłowski TT. ed. *Seed biology*, Vol. III. New York, NY: Academic Press, 145–245.
- Hay FR, Probert RJ. 1995. The effect of different drying conditions and maturity on desiccation tolerance and seed longevity in *Digitalis purpurea* L. *Annals of Botany* 76: 639–647.
- Hay F, Klin J, Probert R. 2006. Can a post-harvest ripening treatment extend the longevity of *Rhododendron* L. seeds? *Scientia Horticulturae* 111: 80–83.
- Hay F, Adams J, Manger K, Probert R. 2008. The use of non-saturated lithium chloride solutions for experimental control of seed water content. *Seed Science and Technology* 36: 737–746.
- Horbowicz M, Obendorf RL. 1994. Seed desiccation tolerance and storability: dependence on flatulence-producing oligosaccharides and cyclitols – review and survey. *Seed Science Research* 4: 385–405.
- Liu K, Eastwood RJ, Flynn S, Turner RM, Stuppy WH. 2008. *Seed information database* (release 7.1). <http://www.kew.org/data/sid>
- Long RL, Panetta FD, Steadman KJ, et al. 2008. Seed persistence in the field may be predicted by laboratory-controlled aging. *Weed Science* 56: 523–528.
- Pagel MD. 1992. A method for the analysis of comparative data. *Journal of Theoretical Biology* 156: 431–442.
- Peco B, Traba J, Levassor C, Sánchez AM, Azcárate FM. 2003. Seed size, shape and persistence in dry Mediterranean grass and scrublands. *Seed Science Research* 13: 87–95.
- Priestley DA. 1986. *Seed aging*. Ithaca, NY: Comstock Publishing Associates.
- Priestley DA, Cullinan VI, Wolfe J. 1985. Differences in seed longevity at the species level. *Plant, Cell & Environment* 8: 557–562.
- Pritchard HW, Dickie JB. 2003. Predicting seed longevity: the use and abuse of seed viability equations. In: Smith RD, Dickie JB, Linington SH, Pritchard HW, Probert RJ. eds. *Seed conservation: turning science into practice*. London: Royal Botanic Gardens, Kew, 655–721.
- Purvis A, Rambaut A. 1995. Comparative analysis by independent contrast (CAIC): an Apple Macintosh application for analysing comparative data. *Computer Applications in the Biosciences* 11: 247–251.
- Purvis A, Gittleman JL, Luh H-K. 1994. Truth or consequences: effects of phylogenetic accuracy on two comparative methods. *Journal of Theoretical Biology* 167: 293–300.
- Rao NK, Hanson J, Dulloo ME, Ghosh K, Nowell D, Larinde M. 2006. *Manual of seed handling in genebanks. Handbooks for genebanks no. 8*. Rome: Bioversity International.
- Roberts EH. 1973. Predicting the storage life of seeds. *Seed Science and Technology* 1: 499–514.
- SCBD (Secretariat of the Convention on Biological Diversity). 2006. *Global strategy for plant conservation*. <http://www.cbd.int/gspc>
- Sokal RR, Rohlf FJ. 1995. *Biometry*, 3rd edn. New York, NY: Freeman and Co.
- Thompson K, Band SR, Hodgson JG. 1993. Seed size and shape predict persistence in soil. *Functional Ecology* 7: 236–241.
- Walters C. 2003. Optimising seed banking procedures. In: Smith RD, Dickie JB, Linington SH, Pritchard HW, Probert RJ. eds. *Seed conservation: turning science into practice*. London: Royal Botanic Gardens, Kew, 723–743.
- Walters C, Wheeler LM, Grotenhuis JM. 2005. Longevity of seeds stored in a genebank: species characteristics. *Seed Science Research* 15: 1–20.

APPENDIX

Details of species used in the study: classification to family level (following APG, 2003), country of origin, time to 50% viability loss (p_{50}) in the ageing conditions (45°C, 60% RH), and germination conditions used. Species in bold were aged at 60°C, 60% RH (p_{50} shown for these is an estimate based on the measured p_{50} at 60°C multiplied by a factor of 10.9; see text for details).

Species name	Family (-aceae)	Country of origin	p_{50} (d)	Germination pre-treatments	Germination conditions (°C)
<i>Acacia etbaica</i> Schweinf.	Fab-	Kenya	90-1	Mechanical scarification with a scalpel	20
<i>Allocauarina acutivalvis</i> (F.Muell.) L.A.S.Johnson	Casuarin-	Australia	202-0		10
<i>Allocauarina campestris</i> (Diels) L.A.S.Johnson	Casuarin-	Australia	221-0		15
<i>Alnus glutinosa</i> (L.) Gaertn.	Betul-	England	10-1		25
<i>Alonsoa acutifolia</i> Ruiz & Pav.	Scrophulari-	Peru	17-1		15
<i>Amaranthus caudatus</i> L.	Amaranth-	England	70-6		20
<i>Anagallis arvensis</i> L.	Primul-	Egypt	140-0		20
<i>Anagallis foemina</i> Mill.	Primul-	England	44-0		25 [†]
<i>Aquilegia formosa</i> Fisch. ex DC.	Ranuncul-	USA	12-7		25
<i>Astronium fraxinifolium</i> Schott	Anacardi-	Brazil	19-0		20
<i>Atriplex bunburyana</i> F.Muell.	Amaranth-	Australia	17-4		15
<i>Atriplex eardleyae</i> Aellen	Amaranth-	Australia	30-0		20
<i>Betula nana</i> L.	Betul-	Scotland	10-6		30
<i>Blackstonia perfoliata</i> (L.) Huds.	Gentian-	England	37-6		20
<i>Brassica napus</i> L.	Brassic-	England	65-1		20
<i>Caiophora lateritia</i> Benth.	Loas-	England	4-3		20
<i>Calceolaria hypericina</i> Poepp. ex Benth.	Scrophulari-	Chile	26-7		15
<i>Calluna vulgaris</i> (L.) Hull	Eric-	England	17-6		35/20* [†]
<i>Calothamnus crassus</i> (Benth.) Hawkeswood	Myrt-	Australia	375-0		15
<i>Calothamnus graniticus</i> Hawkeswood	Myrt-	Australia	256-3		20
<i>Calothamnus robustus</i> Schauer	Myrt-	Australia	374-0		15
<i>Calothamnus rupestris</i> Schauer	Myrt-	Australia	771-0		15
<i>Calotropis procera</i> (Aiton) W.T.Aiton	Asclepiad-	Morocco	18-5		30
<i>Campanula pyramidalis</i> L.	Campanul-	Hrvatska	3-9		20
<i>Campanula rapunculoides</i> L.	Campanul-	Bulgaria	23-0		20
<i>Carex hystericina</i> Muhl. ex Willd.	Cyper-	USA	67-8		25
<i>Centaureum erythraea</i> Rafn	Gentian-	England	36-9		15

Continued

APPENDIX *Continued*

Species name	Family (-aceae)	Country of origin	p_{50} (d)	Germination pre-treatments	Germination conditions (°C)
<i>Cephalaria joppica</i> (Spreng.) Beg.	Dipsac-	Lebanon	128.3		15
<i>Cercidiphyllum japonicum</i> Siebold & Zucc.	Cercidiphyll-	South Korea	0.1		35/20*
<i>Chamaecrista mimosoides</i> (L.) Greene	Fab-	Burkina Faso	63.5	Mechanical scarification with a scalpel	25
<i>Chenopodium auricomum</i> Lindl.	Amaranth-	Australia	36.7		25/10*
<i>Cistus ladanifer</i> L.	Cist-	Spain,	78.2		15
<i>Clarkia botaie</i> (Spach) F.H.Lewis & M.E.Lewis	Onagr-	USA	44.7		15
<i>Clarkia lewisii</i> P.H.Raven & D.R.Parn.	Onagr-	USA	52.8		15
<i>Clarkia rubicunda</i> (Lindl.) F.H.Lewis & M.E.Lewis	Onagr-	USA	49.4		25/10*
<i>Convolvulus crenatifolius</i> Ruiz & Pav.	Convolvul-	Peru	103.4		23/9*
<i>Conyza sumatrensis</i> (Retz.) E.Walker	Aster-	Indonesia, Sulawesi	35.5		20
<i>Cooperia drummondii</i> Herb.	Amaryllid-	USA	8.0		20
<i>Cooperia pedunculata</i> Herb.	Amaryllid-	USA	18.2		20
<i>Cortusa matthioli</i> L.	Primul-	England	3.7		20
<i>Cotula coronopifolia</i> L.	Aster-	New Zealand	105.0		25
<i>Crassula clavata</i> N.E.Br.	Crassul-	South Africa	77.2		20
<i>Cryptantha watsonii</i> (A. Gray) Greene	Boragin-	USA	47.0		10
<i>Cucumis pustulatus</i> Naudin ex Hook. f.	Cucurbit-	Kenya	54.0		35/20*
<i>Cuscuta campestris</i> Yunck.	Convolvul-	France	35.3		25
<i>Cyperus luzulae</i> (L.) Retz.	Cyper-	Peru	26.3		35/20*
<i>Dasyllirion lucidum</i> Rose	Dracaen-	Mexico	73.9		25
<i>Dasyllirion wheeleri</i> S.Watson ex Roth.	Dracaen-	USA	60.3		35/20*
<i>Dipsacus fullonum</i> L.	Dipsac-	England	14.6		15
<i>Dissotis elliotii</i> Gilg	Melastomat-	Burkina Faso	46.5		20
<i>Elymus canadensis</i> L.	Po-	USA	25.4		20
<i>Ephedra fragilis</i> Desf.	Ephedr-	Tunisia	45.9		20
<i>Epilobium ciliatum</i> Raf.	Onagr-	USA	34.6		20
<i>Eucalyptus gracilis</i> F.Muell.	Myrt-	Australia	364.0		20
<i>Ferula communis</i> L.	Api-	Morocco	75.1		10
<i>Ficus exasperata</i> Vahl	Mor-	Malawi	76.6		20
<i>Ficus salicifolia</i> Vahl	Mor-	Oman	33.8		20
<i>Filipendula vulgaris</i> Moench	Ros-	England	16.3		15
<i>Fouquieria splendens</i> Engelm. in Wisl.	Fouquieri-	USA	34.8		15
<i>Galium mollugo</i> L.	Rubi-	Norway	8.3		20
<i>Garidella nigellastrum</i> L.	Ranuncul-	England	28.9		23/9* [†]
<i>Genista umbellata</i> Clos	Fab-	Spain	184.8	Mechanical scarification with a scalpel	15
<i>Gentiana depressa</i> D.Don	Gentian-	Hungary	8.6		20
<i>Gentiana gracilipes</i> Turrill	Gentian-	England	11.0		20
<i>Gentiana purdomii</i> C.Marquand	Gentian-	Finland	8.0		20
<i>Gentiana pyrenaica</i> L.	Gentian-	Germany	13.0		25
<i>Gentiana waltonii</i> Burkill	Ranuncul-	Austria	9.1		15
<i>Geranium lucidum</i> L.	Gerani-	England	113.5		15
<i>Geranium robertianum</i> L.	Gerani-	France	52.1		20
<i>Geum aleppicum</i> Jacq.	Ros-	Canada	27.2		20
<i>Gilia capitata</i> Sims	Polemini-	England	32.2		20
<i>Glaucium corniculatum</i> (L.) Rudolph	Papaver-	Morocco	89.5		35/20* [†]
<i>Gomphocarpus fruticosus</i> (L.) W.T.Aiton	Asclepiad-	Botswana	41.0		25/15*
<i>Habranthus tubispathus</i> (L'Hér.) Traub	Amaryllid-	USA	8.9		20
<i>Hakea candolleana</i> Meisn.	Prote-	Australia	166.0		35/20
<i>Hakea leucoptera</i> R.Br.	Prote-	Australia	14.9		35/20
<i>Hyoscyamus niger</i> L.	Solan-	England	32.0		30 [†]
<i>Hyptis suaveolens</i> (L.) Poit.	Lami-	Nepal	138.8		20
<i>Isopyrum fumarioides</i> L.	Ranuncul-	England	9.8		20
<i>Jasione montana</i> L.	Campanul-	England	10.7		15
<i>Juncus xiphioides</i> E.Mey.	Junc-	USA	59.2		15
<i>Kalanchoe lanceolata</i> (Forssk.) Pers.	Crassul-	Kenya	36.8		25
<i>Kniphofia rooperi</i> (T.Moore) Lem.	Asphodel-	RSA	26.4		25/10*
<i>Leonurus sibiricus</i> L.	Lami-	Argentina	32.2		25/15*
<i>Leptidium montanum</i> Nutt.	Brassic-	USA	22.2		15
<i>Leptospermum ericoides</i> A.Rich.	Myrt-	New Zealand	152.0		25
<i>Leucaena leucocephala</i> (Lam.) de Wit	Fab-	Yemen	75.2	Mechanical scarification with a scalpel	20
<i>Lobelia giberroa</i> Hemsl.	Campanul-	Kenya	7.1		15

Continued

APPENDIX *Continued*

Species name	Family (-aceae)	Country of origin	p_{50} (d)	Germination pre-treatments	Germination conditions (°C)
<i>Ludwigia alternifolia</i> L.	Onagr-	USA	24.1		35/20*
<i>Ludwigia leptocarpa</i> (Nutt.) H.Hara	Onagr-	USA	50.2		35/20*
<i>Lychnis viscaria</i> L.	Caryophyll-	Denmark	6.5		20
<i>Lythrum salicaria</i> L.	Lythr-	England	14.0		23/9* [†]
<i>Meconopsis horridula</i> Hook.f. & Thomson	Papaver-	England	4.8		20
<i>Melaleuca diosmifolia</i> Andrews	Myrt-	Australia	289.0		15
<i>Melaleuca eleuterostachya</i> F.Muell.	Myrt-	Australia	346.0		25
<i>Melaleuca thyoides</i> Turcz.	Myrt-	Australia	420.0		25
<i>Melicytus ramiflorus</i> Forster & Forster f.	Viol-	New Zealand	0.3		15
<i>Mentzelia lindleyi</i> Torr. & A.Gray	Loas-	England	13.7		15
<i>Myrica serrata</i> Lam.	Myric-	Swaziland	11.0		25/10*
<i>Narthecium ossifragum</i> (L.) Huds.	Melanthi-	England	0.1	Warm stratification (28 d at 25°C) followed by cold stratification (56 d at 5°C)	15
<i>Navarretia squarrosa</i> (Eschsch.) Hook. & Arn.	Polemini-	Canada	19.8		20
<i>Nemophila menziesii</i> Hook. & Arn.	Hydrophyll-	USA	51.8		15
<i>Neohymenopogon parasiticus</i> (Wall.) Bennet	Rubi-	Nepal	27.0		25
<i>Nicotiana glutinosa</i> L.	Solan-	England	177.7		20
<i>Nicotiana tabacum</i> L.	Solan-	England	18.5		20
<i>Nolina texana</i> S. Watson	Dracaen-	USA	55.4		15
<i>Oenothera epilobiifolia</i> Kunth	Onagr-	Colombia	71.5		20
<i>Oenothera fallax</i> Renner em Rostanski	Onagr-	France	99.6		25
<i>Oenothera kunthiana</i> (Spach) Munz	Onagr-	Mexico	76.2		15
<i>Oenothera rosea</i> L'Hér. ex Aiton	Onagr-	USA	52.5		20
<i>Oenothera texensis</i> P.H.Raven & D.R.Parn.	Onagr-	USA	37.3		15
<i>Olsynium junceum</i> (E.Mey. ex C.Presl) Goldblatt	Irid-	Chile	53.7		15
<i>Oxychloris scariosa</i> (F.Muell.) Lazarides	Po-	Australia	53.4		20
<i>Paederia grandidieri</i> Drake	Rubi-	Madagascar	12.5		25
<i>Papaver orientale</i> L.	Papaver-	Turkey	15.8		20
<i>Paspalum conjugatum</i> P.J.Bergius	Po-		18.1		25
<i>Phacelia viscida</i> Torr.	Hydrophyll-	USA	47.2		20 [†]
<i>Philadelphus lewisii</i> Pursh	Hydrange-	USA	6.9	Cold stratification (56 d at 5°C)	20
<i>Phycella scarlatina</i> Ravenna	Amarylloid-	Chile	16.7		15
<i>Physalis pubescens</i> L.	Solan-	England	26.5		25
<i>Pieris formosa</i> D.Don	Eric-	England	3.6		20
<i>Pinus maximinoi</i> H.E.Moore	Pin-	Nicaragua	6.7		25/15*
<i>Poa bulbosa</i> L.	Po-	Greece	10.0		15
<i>Podotherca gnaphalioides</i> Graham	Aster-	Australia	43.8		15
<i>Polemonium pauciflorum</i> S.Watson	Polemini-	England	14.0		20/10*
<i>Portulaca oleracea</i> L.	Portul-	Colombia	52.3		25
<i>Primula acaulis</i> (L.) Hill	Primul-	England	21.6		20
<i>Primula cortusoides</i> L.	Primul-	England	10.7		25
<i>Primula magellanica</i> Lehm.	Primul-	Chile	8.0		20/10* [†]
<i>Primula polyneura</i> Franch.	Primul-	England	5.8		20
<i>Primula veris</i> L.	Primul-	England	11.9		20
<i>Pterostyrax hispida</i> Siebold & Zucc.	Stryr-	England	25.1		5
<i>Pycnostachys deflexifolia</i> Baker	Lami-	Kenya	139.6		25
<i>Ranunculus acris</i> L.	Ranuncul-	England	15.2		35/20*
<i>Ranunculus sceleratus</i> L.	Ranuncul-	England	13.7		25/10*
<i>Regelia inops</i> (Schauer) Schauer	Myrt-	Australia	314.0		15
<i>Rhododendron campanulatum</i> D.Don.	Eric-	Nepal	2.2		15
<i>Rhododendron micranthum</i> Turcz.	Eric-	South Korea	4.6		20
<i>Rhododendron mucronulatum</i> Turcz.	Eric-	England	18.8		25
<i>Rhodophiala advena</i> (Ker Gawl.) Traub	Amarylloid-	Chile	23.3		20
<i>Rhodophiala advena</i> (Ker Gawl.) Traub	Amarylloid-	Chile	30.8		15
<i>Rhodophiala bagnoldii</i> (Herb.) Traub	Amarylloid-	Chile	29.7		25
<i>Rhodophiala bagnoldii</i> (Herb.) Traub	Amarylloid-	Chile	12.5		25
<i>Ribes alpinum</i> L.	Grossulari-	England	10.5		15
<i>Ribes sanguineum</i> Pursh	Grossulari-	USA	30.6		25/10*
<i>Rogeria longiflora</i> (Royen) Gay ex Dc.	Pedali-	RSA	189.6		25
<i>Rumex crispus</i> L.	Polygon-	England	38.7		20
<i>Ruschia barnardii</i> L.Bolus	Aizo-	RSA	76.9		20
<i>Sagina procumbens</i> L.	Caryophyll-	England	52.3		20
<i>Salvia disermas</i> L.	Lami-	RSA	59.9		20
<i>Samolus ebracteatus</i> Kunth	Primul-	USA	45.9		20

Continued

APPENDIX *Continued*

Species name	Family (-aceae)	Country of origin	p_{50} (d)	Germination pre-treatments	Germination conditions (°C)
<i>Saponaria officinalis</i> L.	Caryophyll-	France	17.6	Cold stratification (28 d at 5°C)	25/10*
<i>Saxifraga cespitosa</i> L.	Saxifrag-	Wales	6.7		15
<i>Saxifraga petraea</i> L.	Saxifrag-	Italy	2.5		15
<i>Scabiosa prolifera</i> L.	Dipsac-	Portugal	35.1		10
<i>Senna occidentalis</i> (L.) Link	Fab-	Zimbabwe	75.7	Mechanical scarification with a scalpel	20
<i>Silene alba</i> (Mill.) E.H.L.Krause	Caryophyll-	England	65.4		25/10*
<i>Silene colorata</i> Poir.	Caryophyll-	Italy	32.1		20
<i>Silene compacta</i> Fisch. ex Hornem.	Caryophyll-	Greece	51.9		25/10*
<i>Silene conica</i> L.	Caryophyll-	Greece	61.4		20
<i>Silene dioica</i> (L.) Clairv.	Caryophyll-	Sweden	22.3		20
<i>Silene gallica</i> L.	Caryophyll-	Spain	114.7		20
<i>Silene italica</i> (L.) Pers.	Caryophyll-	Greece	40.5		20
<i>Silene mellifera</i> Boiss. & Reut	Caryophyll-	Spain	27.5		15
<i>Silene noctiflora</i> L.	Caryophyll-	Canada	27.4		25/10*
<i>Silene nutans</i> L.	Caryophyll-	Bulgaria	35.5		20
<i>Silene otites</i> (L.) Wibel	Caryophyll-	Germany	9.1		10
<i>Silene vulgaris</i> (Moench) Garcke	Caryophyll-	Germany	43.2		15
<i>Solanum lasiophyllum</i> Humb. & Bonpl. ex Dunal	Solan-	Australia	58.0		35/20*
<i>Solanum panduriforme</i> Dunal	Solan-	Botswana	70.7		35/20*
<i>Solanum quadriculatum</i> F.Muell.	Solan-	Australia	70.0		35/20*
<i>Spergularia marina</i> (L.) Griseb.	Caryophyll-	Argentina	28.8		10
<i>Streptocarpus cyaneus</i> S.Moore	Gesneri-	Swaziland	0.9		20
<i>Swainsona colutooides</i> F.Muell.	Fab-	Australia	12.0		15
<i>Talinum paniculatum</i> (Jacq.) Gaertn.	Portulac-	Argentina	51.1		20
<i>Tetradenia riparia</i> (Hochst.) Codd	Lami-	RSA	7.7		20
<i>Teucrium botrys</i> L.	Lami-	England	41.5		20
<i>Teucrium polium</i> L.	Lami-	Morocco	63.6		15
<i>Thuja plicata</i> Donn ex D.Don	Cupress-		10.7		35/20*
<i>Tofieldia calyculata</i> (L.) Wahlenb.	Melanthi-	Bosnia	2.6	Cold stratification (56 d at 5°C)	20 [†]
<i>Tofieldia pusilla</i> Pers.	Melanthi-	Scotland	11.7	Cold stratification (56 d at 5°C)	20
<i>Torilis radiata</i> Moench	Api-	Lebanon	33.8		20
<i>Triglochin maritima</i> L.	Juncagin-	England	5.6		20
<i>Urtica pilulifera</i> L.	Urtic-	Lebanon	84.4		15
<i>Usteria guineensis</i> Willd.	Logani-	Burkina Faso	40.1		25
<i>Valerianella dentata</i> (L.) Pollich	Valerian-	England	56.6		10 [†]
<i>Veronica verna</i> L.	Scrophulari-	New Zealand	14.0		15
<i>Viola lutea</i> Huds.	Viol-	England	12.2		20
<i>Waitzia nitida</i> (Lindl.) Paul G. Wilson	Aster-	Australia	25.9		15
<i>Weigela subsessilis</i> L.H.Bailey	Caprifoli-	South Korea	8.3	Cold stratification (56 d at 5°C)	33/19*
<i>Withania somnifera</i> (L.) Dunal	Solan-	Kenya	49.5		20
<i>Xerophyta humilis</i> (Baker) T.Durand & Schinz	Vellozi-	Botswana	18.5		25
<i>Xiphidium caeruleum</i> Aubl.	Haemodor-	Ecuador	41.7		25
<i>Yucca angustissima</i> Engelm. ex Trelease	Agav-	USA	42.1		20
<i>Zigadenus elegans</i> Pursh	Melanthi-	Canada	3.8		5

* The elevated temperature was applied for 8 h each day with light provided during the warm phase.

[†] GA₃ at 250 mg L⁻¹ was included in the agar germination medium.